

Brief Report

Increased yield of porcine circovirus-2 by a combined treatment of PK-15 cells with interferon-gamma and inhibitors of endosomal-lysosomal system acidification

G. Misinzo, P. L. Delputte, D. J. Lefebvre, H. J. Nauwynck

Laboratory of Virology, Faculty of Veterinary Medicine, Ghent University, Merelbeke, Belgium

Received 14 June 2007; Accepted 29 September 2007; Published online 17 December 2007

© Springer-Verlag 2007

Summary

Treatment of porcine kidney (PK-15) cells with either interferon-gamma (IFN- γ) or endosomal-lysosomal system acidification inhibitors increases replication of porcine circovirus type 2 (PCV2). In the present study, the effect of a combination of these treatments on the number of infected cells and virus yield was tested. The number of PCV2 (Stoon-1010)-infected PK-15 cells increased in cells treated with ammonium chloride ($445 \pm 39\%$ increase), IFN- γ ($446 \pm 8\%$), ammonium chloride + IFN- γ ($1721 \pm 283\%$), chloroquine diphosphate ($1037 \pm 121\%$), chloroquine diphosphate + IFN- γ ($2199 \pm 255\%$), monensin ($950 \pm 178\%$) and monensin + IFN- γ ($1948 \pm 60\%$). Combined IFN- γ and endosomal-lysosomal system acidification inhibitors increased PCV2 yield by up to 50 times compared to untreated PK-15. This augmented virus replication in PK-15 cells may be helpful in the production of PCV2 vaccines.

*

Correspondence: Hans J. Nauwynck, Laboratory of Virology, Faculty of Veterinary Medicine, Ghent University, Salisburylaan 133, 9820 Merelbeke, Belgium
e-mail: hans.nauwynck@UGent.be

Porcine circovirus type 2 (PCV2) is a member of the genus *Circovirus*, family *Circoviridae* [30]. PCV2 infections are mainly associated with a multifactorial disease in pigs, postweaning multisystemic wasting syndrome (PMWS) [1, 7, 9, 12, 17, 27]. Apart from PMWS, PCV2 has also been associated with various disease syndromes in pigs including porcine dermatitis and nephropathy syndrome [3, 5, 19, 21, 31, 35, 37], congenital tremors [6, 14], reproductive failure [16, 18, 21, 28, 29, 33], exudative epidermitis [38], porcine respiratory disease complex [13], proliferative and necrotizing pneumonia [11] and enteritis [4, 15]. *In vitro*, PCV2 replicates in porcine kidney epithelial cells [2, 20, 25, 32], and PCV2 isolation and production is routinely performed using porcine kidney (PK-15) and swine kidney (SK) epithelial cell lines [2, 32]. When PCV2 replication kinetics were studied in a PK-15 epithelial cell line that was inoculated with the prototype PCV2 strain Stoon-1010 [9], 1.8% of cells were infected with a maximum yield of $3.9 \log_{10}$ TCID₅₀/ml in culture supernatant [22]. This low yield slows down the production of PCV2 vaccines. Therefore, increasing the replication of PCV2 in PK-15 cells will allow vaccine production to be scaled up, making the production more efficient and profitable.

Earlier studies have shown that the number of PCV2-infected cells can be enhanced by treatment of PK-15 cells either with interferon-gamma (IFN- γ) [23] or inhibitors of endosomal-lysosomal system acidification such as ammonium chloride (NH₄Cl), chloroquine diphosphate (CQ) and monensin [25]. In the study by Meerts et al. [23], the effect of IFN- γ treatment of PK-15 cells on the PCV2 yield was also investigated, and it was shown that IFN- γ treatment increased the yield of progeny PCV2 by 20 times. Up till now, neither the effect of treatment of PK-15 cells with the endosomal-lysosomal system acidification inhibitors alone nor the effect of treatment of PK-15 cells with the combination of IFN- γ and inhibitors of the endosomal-lysosomal system acidification on PCV2 yield have been investigated. It was the aim of the present study to investigate the combined effect of treating PK-15 cells with IFN- γ and inhibitors of endosomal-lysosomal system acidification on PCV2 infection and PCV2 yield.

To investigate the effect of IFN- γ and inhibitors of the endosomal-lysosomal system acidification on the number of PCV2-infected cells, 2×10^4 PK-15 cells that were free of porcine circoviruses were seeded per well of a 96-well cell culture plate (Nunc, Roskilde, Denmark). Cells were maintained at 37 °C in a humidified 5% CO₂ incubator in culture medium containing 10% fetal bovine serum (FBS; Invitrogen, Grand Island, U.S.A.), 0.3 mg/ml L-glutamine (BDH Chemicals Ltd., Poole, England), 1% nonessential amino acids (100 \times ; Invitrogen), 1 mM sodium pyruvate (Invitrogen), 100 U/ml penicillin, 0.1 mg/ml streptomycin and 0.1 mg/ml kanamycin in RPMI-1640 (Invitrogen). Recombinant porcine IFN- γ (R & D systems, Abingdon, U.K.) was dissolved and stored as previously described [23]. Cell culture medium with or without 500 U/ml IFN- γ was added to PK-15 cells at 6 h post-seeding. At 24 h post-seeding, cells were washed and inoculated with prototype PCV2 strain Stoon-1010 (5.3 log₁₀ TCID₅₀, 20th passage) at a




Fig. 1. Effect of IFN- γ and inhibitors of endosomal-lysosomal system acidification on PCV2 infection of PK-15 cells. Representative light microscopic images of PK-15 cells treated with or without IFN- γ and/or CQ after performing an IPMA staining to identify PCV2-infected cells. Bar: 200 μ m

multiplicity of infection (moi) of 0.3 at 37 °C. After 1 h, the viral inoculum was washed off and cells were incubated in cell culture medium with or without 25 mM, 125 µM or 6 µM of NH₄Cl, CQ or monensin, respectively. At 24 h post-inoculation (hpi), cell culture medium with or without the endosomal-lysosomal system inhibitors was replaced with fresh cell culture medium without inhibitors of endosomal-lysosomal system acidification. Cells were fixed with methanol at -20 °C for 10 min after the first cycle of PCV2 replication at 36 hpi [22]. PCV2-infected cells were identified by an immunoperoxidase monolayer assay (IPMA) using porcine polyclonal anti-PCV2 antibodies [34] and horseradish peroxidase-conjugated polyclonal rabbit anti-swine immunoglobulins (DakoCytomation, Glostrup, Denmark). Substrate was added to stain-infected cells, which were then counted by examination under a light microscope (Olympus Optical Co., Hamburg, Germany). The number of infected cells per well in untreated PK-15 cells was used as a reference, and all results were expressed as relative percentages to this reference. All experiments were performed three times with each condition in a single experiment in quadruplicate.

Treatment of PK-15 cells either with IFN-γ or inhibitors of endosomal-lysosomal system acidification increased the number of PCV2-infected cells (Figs. 1 and 2), in agreement with previous studies [23, 25]. Combined treatment of PK-15 cells with IFN-γ and inhibitors of endosomal-lysosomal system acidification also increased the number of PCV2-infected cells. The percentages of PCV2-




Fig. 2. The effect of IFN-γ and inhibitors of endosomal-lysosomal system acidification on the number of PCV2-infected PK-15 cells. PK-15 cells were pre-treated with (filled bars) or without (empty bars) IFN-γ before they were inoculated with PCV2. After PCV2 inoculation, cells were incubated with or without inhibitors of endosomal-lysosomal system acidification for 24 h. Cells were fixed at 36 hpi and PCV2-infected cells were counted following IPMA staining. The number of PCV2-infected cells in untreated cell cultures was taken as reference, and all values are relative percentages of this reference

infected PK-15 cells at 36 hpi in untreated and treated PK-15 cells are shown in Table 1. From these results, it can be concluded that inhibiting endosomal-lysosomal system acidification gives an additional enhancement of PCV2 infection in PK-15 cells treated with IFN-γ. This additional effect was possible because IFN-γ and inhibitors of endoso-

Table 1. Total PCV2 yield in untreated PK-15 cells and PK-15 cells treated with IFN-γ, inhibitors of endosomal-lysosomal system acidification, or a combination of IFN-γ and inhibitors of endosomal-lysosomal system acidification

Lysosomotropic agent	IFN-γ (U/ml)	% of PCV2-infected cells at 36 hpi	PCV2 titre log ₁₀ (TCID ₅₀ per 10 ⁵ cells)			
			1 hpi	24 hpi	48 hpi	72 hpi
Control	0	0.38 ± 0.03	1.9	1.5	2.9	3.1
Control	500	1.71 ± 0.12	2.0	2.1	3.4	3.9
NH ₄ Cl	0	1.70 ± 0.06	2.1	1.9	3.0	3.8
NH ₄ Cl	500	6.67 ± 1.57	1.6	3.0	4.0	4.8
CQ	0	3.97 ± 0.31	1.4	1.8	3.3	4.3
CQ	500	8.47 ± 1.23	1.6	2.3	4.0	4.6
Monensin	0	3.61 ± 0.39	1.9	2.3	3.9	3.6
Monensin	500	7.52 ± 1.17	2.0	2.8	4.6	4.8

mal-lysosomal system acidification are known to affect different stages of PCV2 infection. Previous studies have shown that treatment of cells with IFN- γ increases PCV2 infection by increasing internalization of PCV2 particles into the cell [23], while treatment of cells with inhibitors of the endosomal-lysosomal system acidification increases PCV2 infection by increasing disassembly of internalized PCV2 particles within the cell [25].

For virus yield assays, 2×10^5 cells were seeded per well of a 24-well cell culture plate (Nunc). Cells were pre-treated with or without 500 U/ml IFN- γ at 6 h post-seeding. At 24 h post-seeding, cells were washed and inoculated with the prototype PCV2 strain Stoon-1010 ($5.3 \log_{10}$ TCID₅₀, 20th passage) at an moi of 0.3 for 1 h at 37 °C. The viral inoculum was washed off, and cells were further incubated in culture medium with or without 25 mM NH₄Cl, 125 μ M CQ or 6 μ M monensin for 24 h. Then, the culture medium with or without endosomal-lysosomal system acidification inhibitors was replaced with fresh culture medium without inhibitors of endosomal-lysosomal system acidification. All inhibitors of endosomal-lysosomal system acidification were removed at 24 hpi because one of the inhibitors, monensin, is a classical exocytosis blocker [8, 10, 26, 36]. At 1, 24, 48 and 72 hpi, the cell culture supernatant was collected. Cells were subjected to three freeze-thaw cycles. Total virus yield (intra- and extracellular virus titres) was determined on PCV2-negative PK-15 cells by inoculation of a tenfold dilution series on PK-15 cells in quadruplicate, as previously described [24]. After 36 h of cultivation at 37 °C in the presence of 5% CO₂, cells were fixed in methanol at -20 °C for 10 min. Viral antigens were detected using an IPMA as described above, and PCV2 titres were expressed as \log_{10} TCID₅₀ per 10^5 cells.

Treatment of PK-15 cells either with IFN- γ or inhibitors of endosomal-lysosomal system acidification increased PCV2 yield (Table 1). Combined treatment of PK-15 cells with IFN- γ and inhibitors of endosomal-lysosomal system acidification further increased PCV2 yield. PCV2 yield increased gradually under all conditions, with the highest yield observed at 72 hpi. The highest recorded PCV2 yield was $4.8 \log_{10}$ TCID₅₀ per 10^5 treated cells

compared to $3.1 \log_{10}$ TCID₅₀ per 10^5 untreated cells at 72 hpi. This is a 50-fold increase in PCV2 yield. The highest increase in PCV2 infection was observed when PK-15 cells were treated with a combination of IFN- γ and CQ or NH₄Cl.

The results of the present study show that a combined inhibition of endosomal-lysosomal system acidification and IFN- γ treatment of PK-15 cells increases the number of PCV2-infected cells and PCV2 yield. Treatment of cells with a combination of inhibitors of endosomal-lysosomal system acidification and IFN- γ may be of use in increasing the yield of PCV2 for PCV2 vaccine production.

Acknowledgments

The authors are grateful to Carine Boone and Chris Bracke for their skillful technical assistance. This work was funded by the European Union (Sixth Framework Programme, Project No. 513928, coordinated by G. Allan). G.M. was supported with a scholarship for candidates from developing countries from the Special Research Fund (BOF) of Ghent University. P.L.D. was supported with a fellowship (B/06524) from BOF of Ghent University.

References

1. Allan GM, McNeilly F, Kennedy S, Daft B, Clarke EG, Ellis JA, Haines DM, Meehan BM, Adair BM (1998) Isolation of porcine circovirus-like viruses from pigs with a wasting disease in the USA and Europe. *J Vet Diagn Invest* 10: 3–10
2. Allan GM, Mc Neilly F, Meehan BM, Kennedy S, Mackie DP, Ellis JA, Clark EG, Espuna E, Saubi N, Riera P, Botner A, Charreyre CE (1999) Isolation and characterisation of circoviruses from pigs with wasting syndromes in Spain, Denmark and Northern Ireland. *Vet Microbiol* 66: 115–123
3. Allan GM, McNeilly E, Kennedy S, Meehan B, Moffett D, Malone F, Ellis J, Krakowka S (2000) PCV-2-associated PDNS in Northern Ireland in 1990. Porcine dermatitis and nephropathy syndrome. *Vet Rec* 146: 711–712
4. Chae C (2005) A review of porcine circovirus 2-associated syndromes and diseases. *Vet J* 169: 326–336
5. Choi C, Chae C (2001) Colocalization of porcine reproductive and respiratory syndrome virus and porcine circovirus 2 in porcine dermatitis and nephrology syndrome by double-labeling technique. *Vet Pathol* 38: 436–441
6. Choi J, Stevenson GW, Kiupel M, Harrach B, Anothayanontha L, Kanitz CL, Mittal SK (2002)

- Sequence analysis of old and new strains of porcine circovirus associated with congenital tremors in pigs and their comparison with strains involved with post-weaning multisystemic wasting syndrome. *Can J Vet Res* 66: 217–224
7. Clark EG (1997) Post-weaning multisystemic wasting syndrome. *Proc Am Assoc Swine Pract* 28: 490–501
 8. Dinter A, Berger EG (1998) Golgi-disturbing agents. *Histochem Cell Biol* 109: 571–590
 9. Ellis J, Hassard L, Clark E, Harding J, Allan G, Willson P, Strokappe J, Martin K, McNeilly F, Meehan B, Todd D, Haines D (1998) Isolation of circovirus from lesions of pigs with postweaning multisystemic wasting syndrome. *Can Vet J* 39: 44–51
 10. Griffiths G, Quinn P, Warren G (1983) Dissection of the Golgi complex. I. Monensin inhibits the transport of viral membrane proteins from medial to trans Golgi cisternae in baby hamster kidney cells infected with Semliki Forest virus. *J Cell Biol* 96: 835–850
 11. Harding JC (2004) The clinical expression and emergence of porcine circovirus 2. *Vet Microbiol* 98: 131–135
 12. Harding JCS, Clark EG, Strokappe JH, Wilson PI, Ellis JA (1998) Postweaning multisystemic wasting syndrome: epidemiology and clinical presentation. *Swine Health Prod* 6: 249–254
 13. Harms PA, Halbur PG, Sorden SD (2002) Three cases of porcine respiratory disease complex associated with porcine circovirus type 2 infection. *J Swine Health Prod* 10: 27–30
 14. Hines RK, Lukert PD (1994) Porcine circovirus as a cause of congenital tremors in newborn pigs. *Proc. Amer. Assoc. Swine Pract.*, pp 344–345
 15. Kim J, Ha Y, Jung K, Choi C, Chae C (2004) Enteritis associated with porcine circovirus 2 in pigs. *Can J Vet Res* 68: 218–221
 16. Kim J, Jung K, Chae C (2004) Prevalence of porcine circovirus type 2 in aborted fetuses and stillborn piglets. *Vet Rec* 155: 489–492
 17. Kiupel M, Stevenson GW, Mittal SK, Clark EG, Haines DM (1998) Circovirus-like viral associated disease in weaned pigs in Indiana. *Vet Pathol* 35: 303–307
 18. Ladekjaer-Mikkelsen AS, Nielsen J, Storgaard T, Botner A, Allan G, McNeilly F (2001) Transplacental infection with PCV-2 associated with reproductive failure in a gilt. *Vet Rec* 148: 759–760
 19. Larochelle R, Magar R, D’Allaire S (2002) Genetic characterization and phylogenetic analysis of porcine circovirus type 2 (PCV2) strains from cases presenting various clinical conditions. *Virus Res* 90: 101–112
 20. Meehan BM, McNeilly F, Todd D, Kennedy S, Jewhurst VA, Ellis JA, Hassard LE, Clark EG, Haines DM, Allan GM (1998) Characterization of novel circovirus DNAs associated with wasting syndromes in pigs. *J Gen Virol* 79 (Pt 9): 2171–2179
 21. Meehan BM, McNeilly F, McNair I, Walker I, Ellis JA, Krakowka S, Allan GM (2001) Isolation and characterization of porcine circovirus 2 from cases of sow abortion and porcine dermatitis and nephropathy syndrome. *Arch Virol* 146: 835–842
 22. Meerts P, Misinzo G, McNeilly F, Nauwynck HJ (2005) Replication kinetics of different porcine circovirus 2 strains in PK-15 cells, fetal cardiomyocytes and macrophages. *Arch Virol* 150: 427–441
 23. Meerts P, Misinzo G, Nauwynck HJ (2005) Enhancement of porcine circovirus 2 replication in porcine cell lines by IFN-gamma before and after treatment and by IFN-alpha after treatment. *J Interferon Cytokine Res* 25: 684–693
 24. Misinzo G, Delputte PL, Meerts P, Lefebvre DJ, Nauwynck HJ (2006) Porcine circovirus 2 uses heparan sulfate and chondroitin sulfate B glycosaminoglycans as receptors for its attachment to host cells. *J Virol* 80: 3487–3494
 25. Misinzo G, Delputte PL, Nauwynck HJ (2007) Inhibition of endosomal-lysosomal system acidification enhances porcine circovirus 2 infection of porcine epithelial cells. *J Virol*, doi: 10.1128/JVI.01229-07
 26. Mollenhauer HH, Morre DJ, Rowe LD (1990) Alteration of intracellular traffic by monensin: mechanism, specificity and relationship to toxicity. *Biochim Biophys Acta* 1031: 225–246
 27. Nayar GP, Hamel A, Lin L (1997) Detection and characterization of porcine circovirus associated with postweaning multisystemic wasting syndrome in pigs. *Can Vet J* 38: 385–386
 28. O’Connor B, Gauvreau H, West K, Bogdan J, Ayroud M, Clark EG, Konoby C, Allan G, Ellis JA (2001) Multiple porcine circovirus 2-associated abortions and reproductive failure in a multisite swine production unit. *Can Vet J* 42: 551–553
 29. Pensaert MB, Sanchez RE Jr, Ladekjaer-Mikkelsen AS, Allan GM, Nauwynck HJ (2004) Viremia and effect of fetal infection with porcine viruses with special reference to porcine circovirus 2 infection. *Vet Microbiol* 98: 175–183
 30. Pringle CR (1999) Virus taxonomy at The XIth International Congress of Virology, Sydney, Australia. *Arch Virol* 144: 2065–2070
 31. Rosell C, Segales J, Ramos-Vara JA, Folch JM, Rodriguez-Arrioja GM, Duran CO, Balasch M, Plana-Duran J, Domingo M (2000) Identification of porcine circovirus in tissues of pigs with porcine dermatitis and nephropathy syndrome. *Vet Rec* 146: 40–43
 32. Rovira A, Balasch M, Segales J, Garcia L, Plana-Duran J, Rosell C, Ellerbrok H, Mankertz A, Domingo M (2002) Experimental inoculation of conventional pigs with porcine reproductive and respiratory syndrome virus and porcine circovirus 2. *J Virol* 76: 3232–3239
 33. Sanchez RE Jr, Nauwynck HJ, McNeilly F, Allan GM, Pensaert MB (2001) Porcine circovirus 2 infection in

- swine foetuses inoculated at different stages of gestation. *Vet Microbiol* 83: 169–176
34. Sanchez RE Jr, Meerts P, Nauwynck HJ, Pensaert MB (2003) Change of porcine circovirus 2 target cells in pigs during development from fetal to early postnatal life. *Vet Microbiol* 95: 15–25
 35. Saoulidis K, Kyriakis SC, Kennedy S, Lekkas S, Miliotis Ch C, Allan G, Balkamos GC, Papoutsis PA (2002) First report of post-weaning multisystemic wasting syndrome and porcine dermatitis and nephropathy syndrome in pigs in Greece. *J Vet Med B Infect Dis Vet Public Health* 49: 202–205
 36. Tartakoff AM, Vassalli P (1977) Plasma cell immunoglobulin secretion: arrest is accompanied by alterations of the golgi complex. *J Exp Med* 146: 1332–1345
 37. Thomson J, Smith B, Allan G, McNeilly F, McVicar C (2000) PDNS, PMWS and porcine circovirus type 2 in Scotland. Porcine dermatitis and nephropathy syndrome. Post-weaning multisystemic wasting syndrome. *Vet Rec* 146: 651–652
 38. Wattrang E, McNeilly F, Allan GM, Greko C, Fossum C, Wallgren P (2002) Exudative epidermitis and porcine circovirus-2 infection in a Swedish SPF-herd. *Vet Microbiol* 86: 281–293